

ON THE REGULARITY OF CR MAPPINGS IN HIGHER CODIMENSION

A. TUMANOV

Abstract. We give a proof of the regularity of Hölder CR homeomorphisms of strictly pseudo convex CR manifolds of higher codimension.

The purpose of this paper is to complete and simplify the proof of one of the main results of [T] on the regularity of CR mappings of strictly pseudo convex CR manifolds of higher codimension. In [T] we introduce a local theory of extremal discs for such manifolds and apply it to the regularity problem. Theorem 7.1 of [T] and Theorem 1 of this paper are the same result. The proof of Theorem 7.1 relies on Theorem 6.10, whose proof is not given in full detail in [T]. Theorem 6.10 says that the union of certain exceptional discs can cover at most a set of measure zero. Thus, the regularity result is proved in [T] only for manifolds for which the conclusion of Theorem 6.10 is true.

Instead of giving the details of the proof of Theorem 6.10, which appears to be quite involved, we give a direct proof of Theorem 7.1. The proof in this paper does not pass through Theorem 6.10, and therefore Theorem 6.10 is no longer needed for proving the regularity of CR maps.

This paper is written as a continuation of [T] and assumes that the reader is familiar with that paper. We use the same notation and refer to the definitions, statements, and formulas of [T] the way they are numbered there, that is by indicating two numbers separated by a period. We refer to objects from this paper by indicating single numbers.

Thus, we prove the following

Theorem 1. *Let M_1 and M_2 be C^∞ smooth generic strictly pseudoconvex normal (see Definition 6.6) CR manifolds in $\mathbf{C}^N$ with generating Levi forms, and let $F : M_1 \rightarrow M_2$ be a homeomorphism such that both F and F^{-1} are CR and satisfy a Lipschitz condition with some exponent $0 < \alpha < 1$. Then F is C^∞ smooth.*

We refer to [T] for a discussion on this and related results. The hypothesis that M_1 and M_2 are normal can be slightly relaxed. That would require reworking Definition

6.6, Lemma 6.7, and Proposition 6.8 by replacing normal discs by non-defective ones. In Definition 6.6 the matrices $A_1, \dots, A_k$ would be linearly independent as linear operators rather than quadratic forms. However, the improvement would be slight, because the author does not even know any examples of manifolds that are not normal. We leave the details to the reader.

The key idea of the proof of Theorem 1 is contained in Proposition 7.5, which says that F preserves the lifts of extremal discs. One of the assumptions of that statement is that the disc f_2 in the target space is normal. We will prove Proposition 7.5 without that assumption. Then the proof of Theorem 7.1 given in [T] will go through without the use of Theorem 6.10. Thus, we need the following refinement of Proposition 7.5.

Proposition 2. *Let M_1 and M_2 be smooth generic manifolds in $\mathbf{C}^N$, and let $F : M_1 \rightarrow M_2$ be a homeomorphism such that both F and F^{-1} are CR and satisfy a Lipschitz condition with some exponent $0 < \alpha < 1$. Let F_1 be the extension of F to a domain D as in Proposition 7.3. Let f_1 be a small stationary disc attached to M_1 such that $f(\Delta) \subset D$, and let f_1^* be a supporting lift of f_1 . Then the disc $f_2 = F_1 \circ f_1$ is also stationary and $f_2^* = f_1^*(F_1' \circ f_1)^{-1}$ (where f_1^* and f_2^* are considered row vectors) is a lift of f_2 .*

The proof is based on a new extremal property of extremal analytic discs which is more suitable for application to CR mappings than Definition 3.1.

We call p a *real trigonometric polynomial* if it has the form $p(\zeta) = \sum_{j=-m}^m a_j \zeta^j$, where $a_{-j} = \bar{a}_j$. We call a real trigonometric polynomial p *positive* if $p(\zeta) > 0$ for $|\zeta| = 1$.

We put $\Delta_r = \{\zeta \in \mathbf{C} : |\zeta| < r\}$; $\Delta = \Delta_1$.

We put $\text{Res}\phi = \text{Res}(\phi, 0)$, the residue of ϕ at 0.

Definition 3. Let f be an analytic disc attached to a generic manifold $M \subset \mathbf{C}^N$. Let $f^* : \Delta \setminus \{0\} \rightarrow T^*(\mathbf{C}^N)$ be a holomorphic map with a pole of order at most 1 at 0. We say that the pair (f, f^*) has a *special extremal property* (SEP) if there exists $\delta > 0$ such that for every positive trigonometric polynomial p there exists $C \geq 0$ such that for every analytic disc $g : \Delta \rightarrow \mathbf{C}^N$ attached to M such that $\|g - f\|_{C(\bar{\Delta})} < \delta$ we have

$$\text{Re Res}(\zeta^{-1} \langle f^*, g - f \rangle p) + C \|g - f\|_{C(\bar{\Delta}_{1/2})}^2 \geq 0. \quad (1)$$

The above extremal property is close to the one introduced by Definition 3.1 in [T]. In particular, we note (Lemma 4) that stationary discs with supporting lifts have SEP. Conversely, we prove (Proposition 8) that SEP implies that f^* is a lift of f .

In formulating SEP we no longer restrict to the discs g with fixed center $g(0) = f(0)$. This helps prove Proposition 8 in case f is defective; see remark after Lemma 6. The radius $1/2$ plays no special role here. In Definition 3, we could even consider f^* defined only in a neighborhood of 0 and replace $1/2$ by a smaller number. Then SEP would still imply that f^* is a lift of f .

Proof of Proposition 2. Since f_1 has a supporting lift f_1^* , then the pair (f_1, f_1^*) has SEP. Then we prove (Lemma 5) that the pair (f_2, f_2^*) also has SEP. Then by Proposition 8, SEP implies that f_2^* is a lift of f_2 and the proposition follows.

Lemma 4. *Let f be a stationary disc attached to a generic manifold $M \subset \mathbf{C}^N$. Let f^* be a supporting lift of f . Then the pair (f, f^*) has SEP with $C = 0$.*

Proof. For every analytic disc g attached to M (not necessarily close to f), we have $\operatorname{Re}\langle f^*, g - f \rangle \geq 0$ on the unit circle $b\Delta$. Multiplying by a positive trigonometric polynomial p and integrating along the circle we immediately get (1) with $C = 0$. The lemma is proved.

Lemma 5. *Under assumptions of Proposition 2, the pair (f_2, f_2^*) has SEP.*

Proof. For every small disc g_2 attached to M_2 , we put $g_1 = F_2 \circ g_2$, where F_2 is the extension of F^{-1} as in Proposition 7.3. If g_2 is close to f_2 in the sup-norm, then for $\zeta \in \bar{\Delta}_{1/2}$ we have

$$|g_1(\zeta) - f_1(\zeta)| \leq C_1 |g_2(\zeta) - f_2(\zeta)|,$$

where C_1 is the maximum of $\|F_2'\|$, the norm of the derivative of F_2 in a neighborhood of the compact set $f_2(\bar{\Delta}_{1/2})$. Then for $\zeta \in \bar{\Delta}_{1/2}$ we have

$$g_2(\zeta) - f_2(\zeta) = F_1(g_1(\zeta)) - F_1(f_1(\zeta)) = F_1'(f_1(\zeta))(g_1(\zeta) - f_1(\zeta)) + R(\zeta)|g_1(\zeta) - f_1(\zeta)|^2,$$

where $|R(\zeta)| \leq C_2$, and C_2 is the maximum of $\|F_1''\|$ in a neighborhood of the compact set $f_1(\bar{\Delta}_{1/2})$. For every positive trigonometric polynomial p , recalling that $f_2^* = f_1^*(F_1' \circ f_1)^{-1}$, we obtain

$$\begin{aligned} & |\operatorname{Res}(\zeta^{-1}\langle f_2^*, g_2 - f_2 \rangle p) - \operatorname{Res}(\zeta^{-1}\langle f_1^*, g_1 - f_1 \rangle p)| \\ &= \left| \frac{1}{2\pi i} \int_{|\zeta|=1/2} \zeta^{-1} \langle f_2^*(\zeta), R(\zeta)|g_1(\zeta) - f_1(\zeta)|^2 \rangle p(\zeta) d\zeta \right| \\ &\leq C_1^2 C_2 \|p f_2^*\|_{C(b\Delta_{1/2})} \|g_2 - f_2\|_{C(\bar{\Delta}_{1/2})}^2 \end{aligned} \quad (2)$$

Now by Lemma 4, $\operatorname{Re} \operatorname{Res}(\zeta^{-1}\langle f_1^*, g_1 - f_1 \rangle p) \geq 0$, and (2) implies that (f_2, f_2^*) has SEP. The proof is complete.

Lemma 6. Assume a pair (f, f^*) has SEP, where f is a small analytic disc of class $C^\alpha(\bar{\Delta})$ attached to a generic manifold M . Then for every infinitesimal perturbation $\dot{f}$ of f of class $C^\alpha(\bar{\Delta})$, and every real trigonometric polynomial p , we have

$$\operatorname{Re} \operatorname{Res}(\zeta^{-1} \langle f^*, \dot{f} \rangle p) = 0. \quad (3)$$

Proof. We realize $\dot{f}$ by a one parameter family of discs $\zeta \mapsto g(\zeta, t)$ defined for small $t \in \mathbf{R}$ so that $g(\zeta, 0) = f(\zeta)$ and $\frac{d}{dt}|_{t=0} g = \dot{f}$. Plugging g in (1) we get

$$\operatorname{Re} \operatorname{Res}(\zeta^{-1} \langle f^*, g - f \rangle p) + O(t^2) \geq 0.$$

Differentiating at $t = 0$, we obtain (3) for every positive trigonometric polynomial p . Since positive trigonometric polynomials span the set of all real trigonometric polynomials, then the lemma follows.

Remark. We note that if SEP only held for the discs g with fixed center $g(0) = f(0)$, then in the last proof we would have to realize an infinitesimal perturbation $\dot{f}$ with $\dot{f}(0) = 0$ by a family $\zeta \mapsto g(\zeta, t)$ with fixed center $g(0, t) = f(0)$. However, we are able to do it only if f is not defective; see Corollary 2.5.

We observe that for $p \equiv 1$, the condition (3) takes the form

$$\operatorname{Re}(\langle \bar{a}, \dot{f}'(0) \rangle + \langle \bar{b}, \dot{f}(0) \rangle) = 0, \quad (4)$$

where $\bar{a}$ and $\bar{b}$ are the first two Laurent coefficients of f^* , that is

$$f^*(\zeta) = \bar{a}\zeta^{-1} + \bar{b} + \dots \quad (5)$$

The following theorem and its proof are similar to those of Proposition 3.6. The advantage of the new result is that it holds even if the disc f is defective.

Theorem 7. Let f be a small analytic disc of class $C^\alpha(\bar{\Delta})$ attached to a generic manifold $M \subset \mathbf{C}^N$. Assume $a, b \in \mathbf{C}^N$ are such that (4) holds for every infinitesimal perturbation $\dot{f}$ of f of class $C^\alpha(\bar{\Delta})$. Then there exists a unique lift f^* of f of the form (5). Moreover, the correspondence $(f, a, b) \mapsto f^*$ is continuous in the $C^\alpha(b\Delta)$ norm.

Proof. The proof consists of calculations with coordinates. We would very much like to see a natural coordinate-free proof.

We assume that M is given by a local equation $\rho = 0$, where $\rho = x - h(y, w)$, and that G and H are defined by (2.2) and (2.3).

According to the coordinate representation $\mathbf{C}^N = \mathbf{C}_z^k \times \mathbf{C}_w^n$, we put $\dot{f} = (\dot{z}, \dot{w})$, $a = (\lambda', \mu)$, $b = (c', s)$. We also put $\phi = \text{Re}(Gh_w \dot{w})$, $A = 2H(0)^{-1}$, $B = 2(H^{-1})'(0) = -2H(0)^{-1}H'(0)H(0)^{-1}$. Following [T], we use the notation $P_0\psi = (2\pi)^{-1} \int_0^{2\pi} \psi(e^{i\theta}) d\theta$.

By Proposition 2.1, we have

$$\dot{z}(0) = A(P_0\phi + iv), \quad \dot{z}'(0) = B(P_0\phi + iv) + 2AP_0(\bar{\zeta}\phi).$$

We also have $\dot{w}(0) = P_0(\dot{w})$, $\dot{w}'(0) = P_0(\bar{\zeta}\dot{w})$. Plugging the above expressions for $\dot{z}(0)$, $\dot{z}'(0)$, $\dot{w}(0)$, $\dot{w}'(0)$ in (4) we get

$$\text{Re}\{\bar{\lambda}'(B(P_0\phi + iv) + 2AP_0(\bar{\zeta}\phi)) + \bar{\mu}P_0(\bar{\zeta}\dot{w}) + \bar{c}'A(P_0\phi + iv) + \bar{s}P_0\dot{w}\} = 0$$

for every holomorphic vector function $\dot{w}$ and $v \in \mathbf{R}^k$. We put

$$c = \bar{c}'A + \bar{\lambda}'B, \quad \lambda = 2\lambda'\bar{A}. \quad (6)$$

Then (4) takes the form

$$\text{Re}P_0\{(\bar{\lambda}\bar{\zeta} + c)\phi + (\bar{\mu}\bar{\zeta} + \bar{s})\dot{w}\} + \text{Re}(icv) = 0.$$

Since $v \in \mathbf{R}^k$ is arbitrary, then $c \in \mathbf{R}^k$, and recalling the expression of ϕ we obtain

$$\text{Re}P_0\{(\text{Re}(\lambda\zeta + c)Gh_w + \bar{\mu}\bar{\zeta} + \bar{s})\dot{w}\} = 0.$$

Since $\dot{w}$ is arbitrary, then by the moment conditions,

$$\psi = \text{Re}(\lambda\zeta + c)Gh_w + \bar{\mu}\bar{\zeta} + \bar{s}$$

extends holomorphically from $b\Delta$ to Δ , and $\psi(0) = 0$. Put

$$f^*|_{b\Delta} = \text{Re}(\lambda\zeta + c)G\partial\rho. \quad (7)$$

Since ψ is holomorphic, then f^* is a lift of f and has the form

$$f^* = \frac{1}{4}(\lambda\zeta + 2c + \bar{\lambda}\bar{\zeta}^{-1})H dz + (\bar{\mu}\bar{\zeta}^{-1} + \bar{s} - \psi) dw,$$

which immediately implies (5).

The lift f^* defined by (7) is unique because (6) follows by comparing the z -components of (5) and (7), so λ and c are uniquely determined by a and b .

Finally, the mapping $(f, a, b) \mapsto f^*$ is continuous because by (6), λ and c depend continuously on (f, a, b) .

The proof is complete.

Proposition 8. *Assume a pair (f, f^*) has SEP, where f is a small analytic disc of class $C^\alpha(\bar{\Delta})$ attached to a generic manifold M . Then f^* is a lift of f .*

Proof. Let $f^*(\zeta) = \bar{a}\zeta^{-1} + \bar{b} + \dots$. Then by Lemma 6 we have (3), which implies (4). By Theorem 7, there exists a lift $\tilde{f}^*$ of f such that $\tilde{f}^*(\zeta) = \bar{a}\zeta^{-1} + \bar{b} + \dots$. Then $\psi(\zeta) = \zeta^{-1}(\tilde{f}^*(\zeta) - f^*(\zeta))$ is holomorphic in Δ and $\text{Re Res}(\langle \psi, \dot{f} \rangle p) = 0$ for every infinitesimal perturbation $\dot{f}$ and real trigonometric polynomial p . We will show that this implies $\psi \equiv 0$, whence $f^* = \tilde{f}^*$ is a lift of f .

Take $p(\zeta) = c\zeta^m + \bar{c}\zeta^{-m}$, where $c \in \mathbf{C}$ and $m > 0$ is integer. Put $h = \langle \psi, \dot{f} \rangle$. Then $0 = \text{Re Res}(\langle \psi, \dot{f} \rangle p) = \text{Re Res}(hp) = \text{Re}(\bar{c}\text{Res}(h\zeta^{-m}))$ for all $c \in \mathbf{C}$. Then $\text{Res}(h\zeta^{-m}) = 0$ for all integers $m > 0$. Hence $h \equiv 0$.

Now for every $\dot{f}$ we have $\langle \psi, \dot{f} \rangle = 0$. If ψ is not identically equal to zero, then $\psi(\zeta) = \lambda\zeta^m + O(\zeta^{m+1})$, for some integer $m \geq 0$ and $\lambda \neq 0$. Then $\langle \lambda, \dot{f}(0) \rangle = 0$ for all $\dot{f}$. Note the subspace $\{\dot{f}(0)\}$ spans $\mathbf{C}^N$ over $\mathbf{C}$ because the w - and y -components of $\dot{f}(0)$ are arbitrary. Hence $\lambda = 0$ and we come to a contradiction. The proof is complete.

Proof of Theorem 1. We only describe the adjustments that should be made to the proof of Theorem 7.1. As in the latter, we choose a normal extremal disc f_1 attached to M_1 . Actually, we only use that the disc f_1 is not defective, so Theorem 5.5 applies. We don't worry that the disc $f_2 = F_1 \circ f_1$ does not have to be normal, so Theorem 6.10 is no longer used.

Another slight adjustment applies to the proof of the fact that if (g_1, g_1^*) is close to (f_1, f_1^*) , then the lift $g_2^* = g_1^*(F_1' \circ g_1)^{-1}$ of $g_2 = F_1 \circ g_1$ is uniformly bounded in the Hölder norm on $b\Delta$. Now by Theorem 7, $g_2^*(\zeta) = a_2\zeta^{-1} + b_2 + \dots$ is uniquely determined by and continuously depends on (g_2, a_2, b_2) . The latter is uniquely determined by (g_1, a_1, b_1) , where $g_1^*(\zeta) = a_1\zeta^{-1} + b_1 + \dots$. The expressions of a_2 and b_2 in terms of a_1 and b_1 only involve F_1 in a neighborhood of $f_1(0)$. Hence g_2^* is uniformly bounded.

The rest of the proof goes through with no changes.

References

- [T] A. Tumanov, Extremal discs and the regularity of CR mappings in higher codimension, *Amer. J. Math.* **123** (2001), 445–473.

Alexander Tumanov, Department of Mathematics, University of Illinois, Urbana, IL 61801.
E-mail: tumanov@uiuc.edu